

Completion of semirings*

Martin Goldstern

Spring 1985; Summer 2002

Abstract

A semiring can be “completed” (i.e., embedded into a semiring in which all infinite sums are defined and satisfy some reasonable properties) iff this semiring can be naturally partially ordered. This construction is “natural” (a left adjoint to the forgetful functor), and quite straightforward.

Definitions and easy facts

Definition 1. A semiring is a structure $(S, +, \cdot, 0, 1)$ such that $(S, +, 0)$ is a commutative monoid, $(S, \cdot, 1)$ is a monoid, and the two distributive laws $x \cdot (y + z) = x \cdot y + x \cdot z$, $(y + z) \cdot x = y \cdot x + z \cdot x$ hold.

Definition 2. A complete semiring $(S, +, \cdot, 0, 1, \Sigma)$ is a semiring in which for any family $(a_i : i \in I)$ the infinite sum $\sum_{i \in I} a_i$ is defined, and the function Σ satisfies the following:

- If $I = \{j, j'\}$ has two elements, then $\sum_{i \in I} a_i = a_j + a_{j'}$
- If $f : I \rightarrow J$ is a bijection, and $a_i = b_{f(i)}$ for all $i \in I$, then $\sum_{i \in I} a_i = \sum_{j \in J} b_j$
- Whenever $I = \bigcup_{k \in K} J_k$ is a disjoint union of sets, and $b_k := \sum_{j \in J_k} a_j$ for all $k \in K$, then $\sum_{i \in I} a_i = \sum_{k \in K} b_k$.
- $x \cdot (\sum_{i \in I} a_i) = \sum_{i \in I} (x \cdot a_i)$, $(\sum_{i \in I} a_i) \cdot x = \sum_{i \in I} (a_i \cdot x)$.
- If all a_i are $= 0$, then $\sum_{i \in I} a_i = 0$.

Remark 3. If S is a complete semiring, then there are cardinal numbers λ_1 (the “characteristic cardinality” of S , similar to the characteristic of a field) and $\lambda_S \leq \max(\lambda_1, \text{cardinality of } S)$ such that

- For any set I there is a subset $J \subseteq I$ of cardinality $\leq \lambda_1$ such that $\sum_{i \in I} 1 = \sum_{i \in J} 1$
- For any set I there is a subset $J \subseteq I$ of cardinality $\leq \lambda_S$ such that $\sum_{i \in I} a_i = \sum_{i \in J} a_i$

This makes it possible to view the function Σ as a set rather than a proper class.

Definition 4. A complete semiring is called *d-complete* if for all $(a_i : i = 1, 2, 3, \dots)$ we have:

If $a_1 + \dots + a_k = c$ for all k , then $\sum_{i \in \{1, 2, \dots\}} a_i = c$.

*In 1985 I wrote a diploma thesis (in German) on “completion of semirings” at the Institute of Algebra at the Vienna University of Technology. My advisor was Werner Kuich.

I never published my thesis, but some people have expressed an interest in seeing it, and references to it appear in a few published papers or books ([2], [6],[5], [8], [1]). So here is an English summary.

Thus, in a d -complete semiring at least the “discrete convergence” will be respected by countable sums.

$\mathbb{N}^{(\infty)}$ (the natural numbers together with infinity) is a d -complete semiring. If S is a d -complete semiring, and Σ is any “alphabet”, then the set $S\langle\langle\Sigma^*\rangle\rangle$, the formal power series over Σ with coefficients in S , will again naturally form a d -complete semiring.

In [6], the notion of ω -continuity was developed, which is a variant of d -completeness.

Definition 5. An “ordered semiring” is a semiring with a partial order $\leq$ such that the operations $+$ and $\cdot$ are weakly monotone in both arguments, and 0 is the least element. In particular, this implies $a \leq a + x$ for all a, x .

On every semiring (or even any monoid) we can define a transitive reflexive relation $\leq$ by

$$a \leq b \Leftrightarrow \exists x (a + x = b)$$

This relation is a quasiorder, called the “natural quasiorder” on $(S, +)$.

Fact 6. Let $(S, +, \cdot, 0, 1)$ be a semiring. The following are equivalent:

1. There exists an ordered semiring T such that S is isomorphic to a subsemiring of T .
2. “ S can be ordered”, i.e.: there exists a partial order on S making S into an ordered semiring.
3. The natural quasiorder is antisymmetric (i.e., a partial order).
4. For all $a, x, y \in S$: ($a + x + y = a$ implies $a + x = a$).

Definition 7. A complete ordered semiring is called “finitary” if: For any $(a_i : i \in I)$, the sum $\sum_{i \in I} a_i$ is the least upper bound of all finite subsums:

$$\sum_{i \in I} a_i = \sup \left\{ \sum_{i \in F} a_i : F \subseteq I \text{ finite} \right\}$$

Fact 8. Every finitary semiring is d -complete.

The characteristic cardinality of any finitary semiring is at most $\aleph_0$.

Main theorem

Theorem 9. Let $(S, +, \cdot, 0, 1, \leq)$ be an ordered semiring. Then there is a finitary (complete ordered) semiring $\bar{S}$ such that S is a subsemiring of $\bar{S}$ (with the induced order).

Moreover, the construction $S \rightarrow \bar{S}$ is “universal”, that is: For all finitary semirings T and all embeddings $f : S \rightarrow T$ (respecting the semiring operations and the order) there is a unique embedding $\bar{f} : \bar{S} \rightarrow T$ (respecting semiring operations, order, and therefore also $\sum$) such that $\bar{f}$ extends f .

In other words: Let $\mathfrak{D}$ be the category of ordered semirings (where maps have to preserve order and the semiring operations), and let $\mathfrak{F}$ be the category of finitary semirings, $U : \mathfrak{F} \rightarrow \mathfrak{D}$ the forgetful functor. Then the map $S \mapsto \bar{S}$ is a left adjoint to U .

However, see example 13.

Proof. Let S^* be the free (multiplicative) monoid over S (consisting of all formal words with letters in the alphabet S , including the empty word), and let $\mathbb{N}\langle S^* \rangle$ be the set of formal polynomials over the alphabet S with coefficients in $\mathbb{N}$ (i.e., maps from S^* into $\mathbb{N}$ which are 0 except on a finite set), with pointwise addition, and multiplication defined by the Cauchy product.

Let $\mathbb{N}^{(\infty)} := \{0, 1, 2, \dots, \infty\}$, and let $\mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle$ be the set of formal power series over the alphabet S with coefficients in $\mathbb{N}^{(\infty)}$ (i.e., the set of all functions from S^* to $\mathbb{N}^{(\infty)}$, again with addition defined pointwise, and the Cauchy product).

$\mathbb{N}\langle S^* \rangle$ is a semiring. As a set (but not as a semiring), S is naturally embedded in $\mathbb{N}\langle S^* \rangle$ through a map $e : S \rightarrow \mathbb{N}\langle S^* \rangle$. There is a natural semiring homomorphism (the “evaluation map”) $\varphi : \mathbb{N}\langle S^* \rangle \rightarrow S$, satisfying $\varphi(e(s)) = s$ for all $s \in S$.

$\mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle$ is a finitary complete semiring (with the natural order), and $\mathbb{N}\langle S^* \rangle$ is to a subsemiring of $\mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle$. We will find $\bar{S}$ as a homomorphic image of $\mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle$.

$$\begin{array}{ccc} \mathbb{N}\langle S^* \rangle & \subseteq & \mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle \\ e \uparrow & & \downarrow \kappa \\ S & \dots \rightarrow & \bar{S} \end{array}$$

For $r, s \in \mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle$ define $r \lesssim s$ iff

For all $p \in \mathbb{N}\langle S^* \rangle$ with $p \leq_{\mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle} r$ there exists $q \in \mathbb{N}\langle S^* \rangle$, $q \leq_{\mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle} s$, and $\varphi(p) \leq_S \varphi(q)$.

Now check that the relation $r \sim s : \Leftrightarrow r \lesssim s \text{ \& } s \lesssim r$ is a congruence relation on the semiring $\mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle$. Moreover, this congruence relation also respects infinite sums, so

$$\bar{S} := \mathbb{N}^{(\infty)}\langle\langle S^* \rangle\rangle / \sim$$

is a complete semiring. Clearly S is embedded in $\bar{S}$ as an ordered semiring.

It is easy to see that $\bar{S}$ is actually a complete finitary semiring.

For the “moreover” part: note that a partially ordered semiring can be made into a complete *finitary* semiring in at most one way. This fact helps to show that the construction $S \mapsto \bar{S}$ is universal. □

Corollary 10. *Let $(S, +, \cdot, 0, 1)$ be a semiring. The following are equivalent:*

1. *S can be ordered.*
2. *S is a subsemiring of some d -complete semiring.*
3. *S is a subsemiring of some complete finitary semiring.*

Proof. (1) $\Rightarrow$ (3) is proved in theorem 9.

(3) $\Rightarrow$ (2) is trivial.

(2) $\Rightarrow$ (1): Assume that $a + x + y = a$. Then $a = a + (x + y) + (x + y) + (x + y) + \dots = (a + x) + (y + x) + (y + x) + \dots = a + x$. □

My thesis contained also the following remark:

Fact 11. *Let $(S, +, \cdot, 0, 1)$ be a semiring. The following are equivalent:*

1. *S is zero sum free (i.e., $x + y = 0$ implies $x = y = 0$.)*
2. *S is a subsemiring of some complete semiring.*

Proof. (2) $\Rightarrow$ (1): Similar to Corollary 10, (2) $\Rightarrow$ (1).

(1) $\Rightarrow$ (2): Adjoin an element ∞ , and declare $\sum_{i \in I} a_i = \infty$ iff the set $\{i : a_i \neq 0\}$ is infinite, or if for some i we have $a_i = \infty$. □

However, it is not clear that this completion will satisfy the infinite distributivity law.

Examples

1. Let $S = \mathbb{N}$, then $\bar{S} = \mathbb{N}^{(\infty)}$.
2. Let S be the semiring of finite subsets of some set X , together with X itself, with addition=union, multiplication=intersection. Then $\bar{S}$ is the full powerset of X .
3. Let S be the set of finite formal languages $L \subseteq \Sigma^*$ over an alphabet Σ , with $L_1 + L_2 = L_1 \cup L_2$, $L_1 \cdot L_2 = \{v \cdot w : v \in L_1, w \in L_2\}$. Then $\bar{S}$ is the set of all formal languages over Σ^* .
4. Example of a complete but not d-complete semiring: $\{0, \text{finite}, \text{infinite}\}$, with the obvious operations.
5. Example of a d-complete nonfinitary semiring:

$$\{0, \text{finite}, \text{countable}, \text{uncountable}\}$$

with the obvious operations. This semiring has uncountable characteristic cardinality.

Remark 12. *Finitary semirings are d-complete and have characteristic cardinality $\leq \aleph_0$.*

The converse is true for finite semirings: If S is finite, d-complete and has characteristic cardinality $\leq \aleph_0$, then S is finitary.

Example of a d-complete semiring with characteristic cardinality $\aleph_0$ which is not finitary:

$$\{0, 1, 2, \dots, \infty - 2, \infty - 1, \infty\}$$

with the obvious operations, e.g.: $n + (\infty - k) = \infty$ if $n \geq k$, and $\infty - (k - n)$ otherwise

The following example shows that the restriction to *finitary* semirings in theorem 9 is reasonable.

Example 13. *There is no embedding f from $\mathbb{N}$ into a complete semiring $\mathfrak{N}$ such that*

For all complete semirings C and all semiring embeddings $g : \mathbb{N} \rightarrow C$ there is a complete semiring embedding $h : \mathfrak{N} \rightarrow C$ with $h \circ f = g$.

Proof. Use C with a large characteristic cardinality. □

References

- [1] Kazimierz Głazek. *Semirings and Their Applications in Mathematics and Information Sciences*. Kluwer, 2002.
- [2] Jonathan S. Golan. *The theory of semirings with applications in mathematics and theoretical computer science*. Longman Scientific & Technical, Harlow, 1992.
- [3] Jonathan S. Golan. *Semirings and their applications*. Kluwer Academic Publishers, Dordrecht, 1999. Updated and expanded version of *The theory of semirings, with applications to mathematics and theoretical computer science*
- [4] Jonathan S. Golan and Huaxiong Wang. On embedding in complete semirings. *Comm. Algebra*, 24(9):2945–2962, 1996.
- [5] Georg Karner. On limits in complete semirings. *Semigroup Forum*, 45(2):148–165, 1992.
- [6] Werner Kuich. Automata and languages generalized to ω -continuous semirings. *Theoret. Comput. Sci.*, 79(1, (Part A)):137–150, 1991. Algebraic and computing treatment of noncommutative power series (Lille, 1988).

- [7] Werner Kuich. Representations and complete semiring morphisms. *Inform. Process. Lett.*, 56(6):293–298, 1995.
- [8] Werner Kuich. Semirings and formal power series: their relevance to formal languages and automata. In *Handbook of formal languages, Vol. 1*, pages 609–677. Springer, Berlin, 1997.

Martin Goldstern
Algebra, TU Wien
Wiedner Hauptstr 8–10 / 118
A-1040 Wien
Österreich / Austria (Europe)
`goldstern@tuwien.ac.at`
<http://info.tuwien.ac.at/goldstern/>